



PeptideStacks.ca

COMMUNITY-REPORTED RESEARCH SUMMARY

CJC-1295 + Ipamorelin

10 MG and 20 MG combined

Reconstitution examples, component conversion and community-reported usage patterns.

PRODUCT SCOPE

Live PeptideStacks variants: 10 MG and 20 MG combined

Prepared 6 August 2026 | Sources reviewed 2026-08-06

Educational use only - not medical advice

CJC-1295 + Ipamorelin - preparation and dose math

These are calculation examples only. The concentration assumptions below are explicit so syringe units are never separated from vial strength and diluent volume.

10 MG combined vial (5/5 MG)

Add 2.5 mL bacteriostatic water

Concentration: 2 mg/mL of each component

Each U-100 unit = 20 mcg of each

20 MG combined vial (10/10 MG)

Add 2.5 mL bacteriostatic water

Concentration: 4 mg/mL of each component

Each U-100 unit = 40 mcg of each

Conversion table

Each component	10 MG setup	20 MG setup
100 mcg + 100 mcg	5 units	2.5 units
150 mcg + 150 mcg	7.5 units	3.75 units
200 mcg + 200 mcg	10 units	5 units
250 mcg + 250 mcg	12.5 units	6.25 units

CALCULATION CHECK

Always recalculate from the actual vial strength and diluent volume. A U-100 unit is 0.01 mL; it is not a dose by itself.

CJC-1295 + Ipamorelin - community-reported patterns

Anecdotal summary of the cited community discussions. These observations are descriptive, not a recommendation or evidence of safety or efficacy.

Recurring report	Frequency seen	Timing in reports
100/100 mcg	Once daily	Often bedtime, separated from food
100/200 mcg	Once or twice daily	Morning and/or bedtime
200/200 mcg	Once daily, often Mon-Fri	Morning or bedtime

What recurred across reports

- The reviewed reports most often clustered around 100-200 mcg of each component per administration.
- Some users split morning and evening administrations; others moved dosing to morning after sleep disruption.
- Reported food timing and cycle length varied, and no community consensus was evident.

Important limitations

- Posts discussed sleep changes, tingling or numbness, water retention and limited biomarker response; these reports are uncontrolled anecdotes.
- Confirm whether a product contains CJC-1295 with DAC or no DAC. They are not interchangeable, and this document addresses the combined PeptideStacks product only.

Important Medical Disclaimer

EDUCATIONAL INFORMATION ONLY

This document summarizes calculation examples and community discussion about CJC-1295 + Ipamorelin. It is not medical advice, a prescription, a diagnosis, a treatment plan, or a recommendation to use, inject, purchase, compound, or administer any product.

COMMUNITY REPORTS ARE NOT CLINICAL EVIDENCE

Online community posts are self-reported, uncontrolled, may be inaccurate, and cannot establish identity, purity, dosing accuracy, safety, efficacy, or causation. The ranges shown describe what was reported; they do not define an appropriate dose.

PROFESSIONAL DIRECTIONS PREVAIL

Use only under the direction of a licensed healthcare professional and qualified dispensing pharmacy. Follow the supplied concentration, diluent, storage, beyond-use date, injection technique, monitoring and individualized directions.

RECONSTITUTION AND INJECTION RISK

Incorrect concentration, syringe selection, unit conversion, handling, storage or sterile technique can cause unintended dosing, contamination, infection, abscess, tissue injury or allergic reaction.

WHEN TO SEEK URGENT CARE

Stop and seek urgent medical attention for trouble breathing, facial or throat swelling, fainting, chest pain, severe confusion, persistent vomiting, or signs of a serious injection-site infection.