



PeptideStacks.ca

COMMUNITY-REPORTED RESEARCH SUMMARY

TB-500

5 MG

Preparation examples, syringe conversion, community-reported patterns and important limitations.

PRODUCT SCOPE

Live PeptideStacks variants: 5 MG

Prepared 7 August 2026 | Sources reviewed 2026-08-07

Educational use only - not medical advice

TB-500 - preparation and dose math

These are calculation examples only. The concentration assumptions below are explicit so syringe units are never separated from vial strength and diluent volume.

5 MG vial - 2 mL example

Add 2 mL bacteriostatic water

Concentration: 2.5 mg/mL

Each U-100 unit = 25 mcg

5 MG vial - 1 mL example

Add 1 mL bacteriostatic water

Concentration: 5 mg/mL

Each U-100 unit = 50 mcg

Conversion table

Reported amount	5 MG setup	5 MG setup
500 mcg	20 units	10 units
1 mg	40 units	20 units
2 mg	80 units	40 units
5 mg	200 units	100 units

CALCULATION CHECK

Always recalculate from the actual vial strength and diluent volume. A U-100 unit is 0.01 mL; it is not a dose by itself.

TB-500 - community-reported patterns

Anecdotal summary of the cited community discussions. These observations are descriptive, not a recommendation or evidence of safety or efficacy.

Recurring report	Frequency seen	Context in reports
2 mg	Twice weekly	Lower loading reports
2.5-5 mg	Twice weekly	Most recurring loading range
2-5 mg	Every two to four weeks	Maintenance discussions

What recurred across reports

- Community discussions most often centered on soft-tissue recovery, mobility and injury-related goals.
- The amount, frequency, route and cycle length varied substantially between reports.
- Reported outcomes were commonly confounded by diet, training, other products or existing treatment.

Important limitations

- TB-500 human safety and dosing are not established; recovery claims are largely extrapolated from preclinical work.
- Community reports are uncontrolled anecdotes and do not establish an appropriate dose, safety, product identity, purity or efficacy.

! Important Medical Disclaimer

EDUCATIONAL INFORMATION ONLY

This document summarizes calculation examples and community discussion about TB-500. It is not medical advice, a prescription, a diagnosis, a treatment plan, or a recommendation to use, inject, purchase, compound, or administer any product.

COMMUNITY REPORTS ARE NOT CLINICAL EVIDENCE

Online community posts are self-reported, uncontrolled, may be inaccurate, and cannot establish identity, purity, dosing accuracy, safety, efficacy, or causation. The ranges shown describe what was reported; they do not define an appropriate dose.

PROFESSIONAL DIRECTIONS PREVAIL

Use only under the direction of a licensed healthcare professional and qualified dispensing pharmacy. Follow the supplied concentration, diluent, storage, beyond-use date, injection technique, monitoring and individualized directions.

RECONSTITUTION AND INJECTION RISK

Incorrect concentration, syringe selection, unit conversion, handling, storage or sterile technique can cause unintended dosing, contamination, infection, abscess, tissue injury or allergic reaction.

WHEN TO SEEK URGENT CARE

Stop and seek urgent medical attention for trouble breathing, facial or throat swelling, fainting, chest pain, severe confusion, persistent vomiting, or signs of a serious injection-site infection.

SOURCE BASIS

PubMed literature search: TB-500 - <https://pubmed.ncbi.nlm.nih.gov/?term=TB-500> - retrieved 2026-08-07